

INDIRECT EVIDENCE FOR AN ACTIVE RADIO PULSAR IN THE QUIESCENT STATE OF THE TRANSIENT MS PULSAR SAX J1808.4–3658

SERGIO CAMPANA¹, PAOLO D’AVANZO¹, JORGE CASARES², STEFANO COVINO¹, GIANLUCA ISRAEL³, GIANNI MARCONI^{3,4}, ROB HYNES⁵, PHIL CHARLES⁶, LUIGI STELLA³
 campana@merate.mi.astro.it

Draft version February 2, 2008

ABSTRACT

Millisecond radio pulsars are neutron stars that have been spun-up by the transfer of angular momentum during the low-mass X-ray binary phase. The transition from an accretion-powered to a rotation-powered pulsar takes place on evolutionary timescales at the end of the accretion process, however it may also occur sporadically in systems undergoing transient X-ray activity. We have obtained the first optical spectrum of the low mass transient X-ray pulsar SAX J1808.4–3658 in quiescence. Similar to the black widow millisecond pulsar B1957+20, this X-ray pulsar shows a large optical modulation at the orbital period due to an irradiated companion star. Using the brightness of the companion star as a bolometer, we conclude that a very high irradiating luminosity, a factor of ~ 100 larger than directly observed, must be present in the system. This most likely derives from a rotation-powered neutron star that resumes activity during quiescence.

Subject headings: accretion, accretion disks — binaries: close — stars: neutron

1. INTRODUCTION

SAX J1808.4–3658 was the first-discovered low mass X-ray binary (LMXB) transient showing coherent pulsations. This confirmed unambiguously the long-sought connection between LMXBs and millisecond radio pulsars (Bhattacharya & van den Hevel 1991; Tauris & van den Hevel 2004). The detection of coherent X-ray pulsations during outbursts testifies that the neutron star possesses a magnetic field of $B \sim 10^8 - 10^9$ G, sufficient for a small magnetosphere to form (Psaltis & Chakrabarty 1999; Menna et al. 2003; Di Salvo & Burderi 2003). Unlike persistent LMXBs, SAX J1808.4–3658 is a transient system, i.e. it is active in X-rays only for short intervals lasting a few months (outbursts) followed by quiescent periods of years.

During quiescence LMXB transients are very faint in X-rays (5–6 orders of magnitude less than in outburst) usually with luminosities of $10^{32} - 10^{33}$ erg s^{−1}. Transient systems, therefore, represent a unique laboratory for the study of compact objects in accretion regimes that are inaccessible to persistent sources. Most neutron star transients are characterized by a quiescent X-ray spectrum consisting of a soft component, usually ascribed to the cooling neutron star which has been heated during outbursts (Brown et al. 1998; Rutledge et al. 1999), and a hard power-law tail (with photon index $\Gamma \sim 1 - 2$) the nature of which is still debated (Campana & Stella 2000). Several mechanisms have been put forward to explain these spectral components, ranging from accretion disks (in different flavours such as advection-dominated or convection-dominated disks, disks stopping at the magnetosphere, etc.; Narayan et al. 1997; Blandford & Begelman 1999; Igumenshchev et al. 2003) to emission from

the interaction between the relativistic wind from a re-activated radio pulsar with matter outflowing from the companion star (Stella et al. 1994; Campana et al. 1998b). XMM-Newton carried out the first detailed study of SAX J1808.4–3658 in quiescence: SAX J1808.4–3658 was dimmer than any other neutron star transient (with a 0.5–10 keV luminosity of 5×10^{31} erg s^{−1} at a distance of 2.5 kpc, in’t Zand et al. 2001) and exhibited only a hard power-law component ($\Gamma = 1.4^{+0.6}_{-0.3}$, 90% confidence level) in its quiescent X-ray spectrum (Campana et al. 2002). These results were confirmed by a subsequent XMM-Newton observation (Wijnands 2002).

In the optical, SAX J1808.4–3658 is dim during quiescence (mean $R \sim 20.9 \pm 0.1$; Homer et al. 2001) and brightens considerably in outbursts ($R \sim 16.2 \pm 0.2$; Wang et al. 2001). The optical light curve in outburst and quiescence is modulated at the orbital period and it is in anti-phase with the X-ray light curve, likely indicating that irradiation of the companion star plays a crucial role in spite of the low X-ray luminosity. This is unlike other quiescent transients. The mass function derived from X-ray data ($4 \times 10^{-5} M_{\odot}$, Chakrabarty & Morgan 1998) and the requirement that the companion fills its Roche lobe led to the conclusion that it must be a rather low mass star, possibly a brown dwarf (Bildsten & Chakabarty 2001). A white dwarf companion is ruled out because it would not fill its Roche lobe. Homer et al. (2001) proposed that the bulk of the optical emission in quiescence arises from the internal energy release of a remnant disc and the orbital modulation from the varying contribution of the heated face of the companion star. Burderi et al. (2003) noted that the required irradiating luminosity needed to match

¹ INAF-Osservatorio Astronomico di Brera, Via Bianchi 46, I-23807, Merate (LC), Italy

² Instituto de Astrofísica de Canarias, 38200 La Laguna, Tenerife, Spain

³ INAF-Osservatorio Astronomico di Roma, Via Frascati 33, I-00040, Roma, Italy

⁴ European Southern Observatory, St. Alonso de Córdova 3107, Vitacura, Casilla 19001, Santiago 19, Chile

⁵ Astronomy Department, University of Texas at Austin, 1 University Station C1400, Austin, TX 78712

⁶ Department of Physics and Astronomy, University of Southampton, Southampton SO17 1BJ, UK

the optical flux, however, is a factor 10–50 higher than the quiescent X-ray luminosity of SAX J1808.4–3658. These authors proposed an alternative scenario, in which the irradiation is due to the rotational energy emitted in the form of a relativistic particle wind from the fast spinning neutron star, which switched to the rotation-powered regime during quiescence. Their results are in agreement with the weak constraints from the optical magnitudes by Homer et al. (2001).

2. DATA ANALYSIS

Here we present indirect evidence for an active radio pulsar in SAX J1808.4–3658 during quiescence with the first optical spectra and I band photometry of SAX J1808.4–3658. These data were obtained with the ESO-VLT (UT4 Yepun) during two half nights on July 12–13 2002. We carried out I band photometry with 3 min exposures with FORS2 (pixel size of $0''.126/\text{pixel}$ and a field of view of $6'.8 \times 6'.8$) over one orbital period. Spectroscopy of the same target was performed using the low resolution grism 600RI (centered at $6780 \text{ \AA}$ with a resolution of $55 \text{ \AA}/\text{mm}$) covering $5120\text{--}8450 \text{ \AA}$ and a $1''$ slit with 3 min spectra over four orbital periods. Data reduction was done in MIDAS to remove the bias level, flat-field.

The region around the optical companion of SAX J1808.4–3658 is crowded and poor seeing conditions complicated the analysis (varying between $1.5\text{--}2''$). We take advantage of previous ESO-VLT images (obtained during quiescence in 1999 with seeing $\sim 0.5''$) to de-blend our data (these magnitudes were $V = 21.82 \pm 0.03$, $R = 21.63 \pm 0.04$ and $I = 21.08 \pm 0.04$). Our I photometry shows a dimmer source ($I = 21.5 \pm 0.1$) and a clear modulation at the 2.01 hr orbital period with a semi-amplitude $0.2 \pm 0.04 \text{ mag}$ (65% in flux, Fig. 2). We also re-calculated the modulation semi-amplitude in V (0.13 ± 0.06) and R (0.39 ± 0.09) using the Homer et al. (2001) data. The I -band light curve shows a clear maximum at phase 0.52 ± 0.05 and a single minimum at phase 0.02 ± 0.05 (based on the precise X-ray ephemerides Chakrabarty & Morgan 1998). This is a clear indication of emission from an irradiated companion (e.g. Charles & Coe 2004) and it argues against emission from the impact point between the gas stream from the companion and an accretion disk (the hot spot) since this has maximum at phase $0.8\text{--}0.9$.

We obtained spectra at 3-min intervals over four orbital periods. We selected spectra taken with seeing better than $1.6''$ (due to poor seeing conditions), collecting a total of 51 min of good data. Wavelength calibrations used HeArNe arc lamp observations. Second-order flexure effects were corrected using night-sky emission lines. This correction was always $< 0.3 \text{ \AA}$. Spectra were corrected for slit-losses according to Diego (1985). We also account for the contaminating stars, estimating their relative contribution in a $1''$ slit on the good-seeing VLT images and interpolating to the spectral range. Errors were tracked along these processes resulting in a 0.1 mag error. A weak $H\alpha$ emission line is visible in the spectrum (equivalent width $EW = 10.3 \pm 3.7 \text{ \AA}$, 68% confidence level, and $FWHM = 44.0 \pm 6.3 \text{ \AA}$, see insert in Fig. 1).

3. MODELLING THE DATA

We first checked that at the time of our observations the source was in quiescence. A Rossi X-ray Timing Explorer (RXTE) pointed observation, performed 15 days before the optical observations, provided a 3σ upper limit of $\sim 10^{34} \text{ erg s}^{-1}$ (0.5–10 keV). Similar limits are provided by the Galactic bulge scan carried out with RXTE (nearest observation on Jul 10, 2002; Swank et al. 2002). While these limits are not particularly constraining, they are sufficient to exclude that the source was in outburst (note that an outburst started three months later Wijngaards et al. 2003).

Since SAX J1808.4–3658 was in quiescence, what is the cause of the optical emission? Likely candidates are emission from the companion star and/or the disk. In order to fit within the Roche lobe of a 2.01 hr binary the companion mass has to be less than $0.17 M_{\odot}$. In the model of Bildsten & Chakrabarty (2001) the most likely companion is a $0.05 M_{\odot}$ brown dwarf bloated by irradiation to fill its Roche lobe ($0.13 R_{\odot}$). The maximum intrinsic optical luminosity from the companion for any of the models by Bildsten & Chakrabarty (2001) is $\sim 3 \times 10^{31} \text{ erg s}^{-1}$ (corresponding to a star temperature of 4800 K for a distance of 2.5 kpc). This is too low a luminosity to account for the observed optical flux, which is a factor of > 10 brighter. We therefore turn to the accretion disk as a possible source. Assuming that the quiescent X-ray luminosity is powered by accretion we can infer the expected mass inflow rate and derive the corresponding optical luminosity (including irradiation), which fails to account for what we see by more than a factor of 100. Disk models may be envisaged with a much higher mass accretion rate together with a truncation radius (fine) tuned to avoid optical and soft X-ray violation of observed data. However, this kind of models still require some additional ingredient to explain the large optical phase modulation. The emission from the pulsar could itself extend to the optical and extrapolation of the power law X-ray flux (assuming the XMM-Newton observation found the X-rays in a similar state) could account for about 30% of the optical luminosity. But this also could not explain the observed orbital phase modulation.

It is instructive to compare the properties of SAX J1808.4–3658 with those of the *black widow* pulsar PSR B1957+20 (Fruchter et al. 1988), which consists of a 1.6 ms radio pulsar irradiating its white dwarf companion (orbital period of 9.16 hr) with a rotational energy of $10^{35} \text{ erg s}^{-1}$. X- and γ -rays are generated in an inter-binary shock front, which causes ablation and heating of the companion (Phynney et al. 1988; Arons & Tavani 1994). An X-ray nebula has recently been revealed around PSR B1957+20 confirming this scenario (Stappers et al. 2003), and the orbital modulation is large with an R -band semi-amplitude $> 4 \text{ mag}$ (Callanan et al. 1995). A similar system is the eclipsing millisecond radio pulsar PSR J2051–0827, consisting of a 4.5 ms pulsar orbiting its very low mass companion ($\sim 0.03 M_{\odot}$) every 2.4 hr (Stappers et al. 1996). Radio eclipses as well as a $\sim 3.3 \text{ mag}$ optical modulation have been observed in PSR J2051–0827 (Stappers et al. 2001), however no X-ray observations are available.

Inspired by this analogy and following Burderi et al. (2003), we now attempt to account for the optical and

X-ray spectra (even if not close in time) as well as the V , R and I modulations of SAX J1808.4–3658 with an irradiated star plus the contribution of the shock front. We fit the data by using the irradiating luminosity (L_{irr}), the fractional luminosity difference between the heated and the cold face of the companion (f) and interstellar absorption (A_V) as free parameters (e.g. Chakrabarty 1998). We obtain a good fit to all the available data (reduced $\chi^2 = 0.7$ with 57 degrees of freedom⁷, see Fig. 1). In particular, the required irradiating luminosity is $L_{\text{irr}} = (4_{-1}^{+3}) \times 10^{33} \text{ erg s}^{-1}$ (90% confidence level for three free parameters, i.e. $\Delta\chi^2 = 6.3$). The best fit fraction is $f = 0.65 \pm 0.10$ resulting in temperature difference at the two faces of the companion star of about $1000 \pm 300 \text{ K}$. This temperature difference is similar to the one observed in PSR J2051–0827. The amplitude modulation of SAX J1808.4–3658 is instead smaller than in the case of PSR J2051–0827 and PSR B1957+20. This could be due to some remaining contamination due to our de-blending process, or it could be an inclination effect. For the most strongly modulated PSR J1957+20, the inclination is $\sim 70^\circ$. PSR J2051–0827 has an inclination of $\sim 40^\circ$.

The estimated absorption is $A_V = 1.0 \pm 0.5$, in line with previous estimates. Optical emission from the shock front accounts for about 15% of the total emission⁸. In the fit we assumed that all the irradiating luminosity is re-emitted by the star. Therefore the irradiating luminosity we derived represents only a lower limit since a relativistic particle wind could have rather different effects from those of X-ray irradiation in terms of effective albedo. Moreover, we note that evidence of this large irradiating luminosity comes from the equivalent width of the $H\alpha$ line. If the line comes from reprocessing, one can roughly infer an irradiating luminosity of $\gtrsim 3 \times 10^{33} \text{ erg s}^{-1}$. This can be estimated by taking the line flux and increasing

it by the fraction of emitted flux intercepted by the star ($R_*^2/4a^2$, with R_* the companion star radius taken equal to the Roche lobe radius and a the orbital separation) and by the fraction of energy re-emitted in $H\alpha$ (e.g. Hynes et al. 2002). Here we assumed a conservative value of 0.3 (see the discussion in Hynes et al. 2002).

4. CONCLUSIONS

The required irradiating luminosity is in all cases large ($\sim 4 \times 10^{33} \text{ erg s}^{-1}$), indeed much larger than the observed X-ray luminosity in quiescence; neither accretion-driven X rays, nor the intrinsic luminosity of the companion star or disk are able to account for it (see also Burderi et al. 2003). The only source of energy available within the system is then the rotational energy of the neutron star. In order to have such a large spin-down luminosity one needs a neutron star magnetic field $\bar{B} \gtrsim 6 \times 10^7 \text{ G}$ (the companion star albedo might be larger than zero and the neutron star emission may also be partially beamed; this value is smaller than the one estimated by Burderi et al. 2003, since they did not account for the contamination from nearby stars). The required magnetic field is well in the range inferred from X-ray observations during the outbursts (see above). Prospects for directly observing SAX J1808.4–3658 pulsing in the X-ray band are rather hard since with XMM-Newton we collected less than 300 counts in 30 ks (Campana et al. 2002). Given the crowding around SAX J1808.4–3658 in the optical band, it would be difficult to detect an $H\alpha$ nebula like that around PSR B1957+20. On the other hand a search in the radio band looking for a millisecond radio pulsar would require searches at high frequencies to overcome the effects of free-free absorption (Campana et al. 1998a; Ergma & Antipova 1999; Burgay et al. 2003; Burderi et al. 2003) and a favorable orientation of the radio beam.

REFERENCES

- Arons, J., Tavani, M. 1994, ApJS, 90, 797
 Bhattacharya, D., van den Heuvel, E. P. J. 1991, Phys. Rep. 203, 1
 Bildsten, L., Chakabarty, D. 2001, ApJ, 557, 292
 Blandford, R. D., Begelman, M. C. 1999, MNRAS, 303, L1
 Brown, E. F., Bildsten, L., Rutledge, R. E. 1998, ApJ, 504, L95
 Burderi, L., et al. 2003, A&A, 404, L43
 Burgay, M., et al. 2003, ApJ, 589, 902
 Callanan, P. J., van Paradijs, J., Rengelink, R. 1995, ApJ, 439, 928
 Campana, S., Colpi, M., Mereghetti, S., Stella, L., Tavani, M. 1998a, A&A Rev., 8, 279
 Campana, S., Stella, L. 2000, ApJ, 541, 849
 Campana, S., et al. 1998b, ApJ, 499, L65
 Campana, S., et al. 2002, ApJ, 575, L15
 Chakrabarty, D. 1998, ApJ, 492, 342
 Chakrabarty, D., Morgan, E. H. 1998, Nat., 394, 364
 Charles, P. A., Coe, M. J. 2004, in *Compact Stellar X-Ray Sources*, eds. W.H.G. Lewin and M. van der Klis, (Cambridge University Press), in press
 Diego, F. 1985, PASP, 97, 1209
 Di Salvo, T., Burderi, L. 2003, A&A 397, 723
 Ergma, E., Antipova, J. 1999, A&A, 343, L45
 Fitzpatrick, E. L. 1999, PASP 111, 63
 Fruchter, A. S., Stinebring, D. R., Taylor, J. H. 1988, Nat., 333, 237
 Homer, L., Charles, P. A., Chakrabarty, D., Zyl, L. 2001, MNRAS, 325, 1471
 Hynes, R. I., et al. 2002, MNRAS, 330, 1009
 Igumenshchev, I. V., Narayan, R., M. A. Abramowicz, M. A. 2003, ApJ, 592, 1042
 in't Zand, J. J. M., et al. 2001, A&A, 372, 916
 Menna, M. T., Burderi, L., Stella, L., Robba, N., van der Klis, M. 2003, ApJ, 589, 503
 Narayan, R., Garcia, M. R., McClintock, J. E. 1997, ApJ, 478, L79
 Phinney, E. S., Evans, C. R., Blandford, R. D., Kulkarni, S. R. 1988, Nat., 333, 832
 Psaltis, D., Chakrabarty, D. 1999, ApJ, 521, 332
 Rutledge, R. E., et al. 1999, ApJ, 514, 945
 Stappers, B. W. et al. 1996, ApJ, 465, L119
 Stappers, B. W. et al. 2001, ApJ, 548, L183
 Stappers, B. W. et al. 2003, Science, 299, 1372
 Stella, L., et al. 1994, ApJ, 423, L47
 Swank, J., Markwardt, C. 2002, in *X-ray Binaries in the Chandra and XMM-Newton Era* <http://asc.harvard.edu/xrbconf/proceedings.html>
 Tauris, T. M., van den Heuvel, E. P. J. 2004, in *Compact Stellar X-Ray Sources*, eds. W.H.G. Lewin and M. van der Klis, (Cambridge University Press), in press
 Wang, Z., et al. 2001, ApJ, 563, L61
 Wijnands, R. 2003, ApJ, 588, 425
 Wijnands, R., et al. 2003, Nat., 424, 44

⁷ The conversion efficiency of rotational energy into X-rays for SAX J1808.4–3658 is $\lesssim 10^{-3}$, a factor of 10 higher than PSR 1957+20 (Stappers et al. 2003), as expected due to geometrical reasons.

⁸ We fit our data with an irradiated star plus the extrapolation to the optical of the X-ray tail. The presence of a disk is not required by our fit. We tried in any case to fit with an irradiated star plus a disk model finding similar results for the star parameters and a large disk inner radius ($\gtrsim 10^9 \text{ cm}$).




Fig. 1. Central panel: X-ray to optical (VLT) spectrum of SAX J1808.4–3658. The data are corrected for absorption ($A_V = 1.0$) using standard extinction curve (Fitzpatrick 1999). Dots shows the XMM-Newton (X-rays) and VLT (optical spectrum plus I photometry) data. The continuous (blue) line shows the overall best fit for an irradiating luminosity of $4 \times 10^{33} \text{ erg s}^{-1}$. The (green) dotted line shows the contribution from shock emission between the relativistic pulsar wind and matter outflowing from the companion. The power law tail has a photon index of $\Gamma = 2.0$, this provides a slightly better fit to the data with respect to a $\Gamma = 1.5$ model (in that case the required irradiating luminosity is higher). The lower continuous (red) line shows the mean contribution from the bloated ($0.13 R_\odot$) companion star. This spectrum together with the optical V , R and I modulation has been fitted with an irradiated star plus shock emission model. Free parameters were the ratio between the ‘hot’ and the ‘cold’ star surface and absorption. A downhill simplex method for the search of the minimum has been adopted. Errors were computed in the 3-dimension space with $\Delta\chi^2 = 6.3$. Upper insert: Optical spectrum with the best fit model spectrum (upper continuous [blue] line). The mean star spectrum (lower continuous [red] line) together with the minimum and maximum spectra (dotted [green] lines) are also shown. Lower insert: particular of the broad ($FWHM = 44.0 \pm 6.3 \text{ \AA}$) $H\alpha$ line. The line equivalent width is $EW = 10.3 \pm 3.7 \text{ \AA}$.




Fig. 2. VLT *I* band light curve (lower panel) plus *V* (upper panel) and *R* (middle panel) taken from Homer et al. (2001) deconvolved light curves of SAX J1808.4–3658. Filled dots represent the data and the blue-cyan heavy line is the best-fit model for the *V* data, the orange line the model for the *R* data and the red line for the *I* data. Phases are from the X-ray ephemeris, i.e. the mean orbital longitude as derived from the X-ray pulse arrival time delays.